

## REPORT DOCUMENTATION PAGE

Estimated time to complete: 1 hour. This does not include the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and sending the comments regarding this burden estimate or any other aspect of the collection of information. Send comments regarding this burden estimate or any other aspect of the collection of information to: Director, Defense Technical Information Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1200, Arlington, VA 22202-1860, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.

1 AGENCY USE ONLY (Leave Blank) 2 REPORT DATE 3 REPORT TYPE AND DATES COVERED

## 4 TITLE AND SUBTITLE

Growth of Third-Harmonic Signal in Optical Glass Fibre

## 5 FUNDING NUMBERS

TA2301-Y1-72

## 6 AUTHOR(S)

Ulf Österberg

## 7 PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Frank J. Seiler Research Laboratory  
USAF Academy CO 80840-6528

## 8 PERFORMING ORGANIZATION REPORT NUMBER

FSJRL-JR-90-0002

## 9. SPONSORING MONITORING AGENCY NAME(S) AND ADDRESS(ES)

N/A

## 10 SPONSORING MONITORING AGENCY REPORT NUMBER

N/A

## 11. SUPPLEMENTARY NOTES

N/A

## 12a. DISTRIBUTION AVAILABILITY STATEMENT

Distribution Unlimited

## 12b. DISTRIBUTION CODE

N/A

## 13. ABSTRACT (Maximum 200 words)

Rapid growth with time of the third-harmonic signal in optical glass fibres, illuminated with 10kW peak power pulses from a Nd:YAG laser, has been observed. Broadband fluorescence from the third-harmonic signal around 420 nm was also detected. *Keywords: JSI*

**S DTIC  
ELECTE  
MAR 26 1990  
E<sup>o</sup> D**



Accession For NTIS GRAIL DTIC TAB Unannounced Justification	By Distribution/ Availability Codes Avail and/or Special
	Dist A-120

## 14. SUBJECT TERMS

Optical fibres, nonlinear optics, optics, glass

(A/w)

## 15. NUMBER OF PAGES

2

## 16. PRICE CODE

## 17. SECURITY CLASSIFICATION OF REPORT

UNCLASSIFIED

## 18. SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

## 19. SECURITY CLASSIFICATION OF ABSTRACT

UNCLASSIFIED

## 20. LIMITATION OF ABSTRACT

NONE

## GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to stay within the lines to meet optical scanning requirements.

**Block 1. Agency Use Only (Leave blank)**

**Block 2. Report Date** Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

**Block 3. Type of Report and Dates Covered** State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88)

**Block 4. Title and Subtitle** A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

**Block 5. Funding Numbers** To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit
	Accession No.

**Block 6. Author(s)** Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

**Block 7. Performing Organization Name(s) and Address(es)** Self-explanatory.

**Block 8. Performing Organization Report Number** Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

**Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es)** Self-explanatory.

**Block 10. Sponsoring/Monitoring Agency Report Number (If known)**

**Block 11. Supplementary Notes** Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

**Block 12a. Distribution/Availability Statement**

Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

**Block 12b. Distribution Code**

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank.

NTIS - Leave blank.

**Block 13. Abstract** Include a brief (Maximum 200 words) factual summary of the most significant information contained in the report.

**Block 14. Subject Terms** Keywords or phrases identifying major subjects in the report.

**Block 15. Number of Pages** Enter the total number of pages.

**Block 16. Price Code** Enter appropriate price code (NTIS only).

**Blocks 17. - 19. Security Classifications** Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

**Block 20. Limitation of Abstract** This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

## GROWTH OF THIRD-HARMONIC SIGNAL IN OPTICAL GLASS FIBRE

*Indexing terms: Optical fibres, Nonlinear optics, Optics, Glass*

Rapid growth with time of the third-harmonic signal in optical glass fibres, illuminated with 10kW peak power pulses from a Nd: YAG laser, has been observed. Broadband fluorescence from the third-harmonic signal around 420 nm was also detected.

Efficient second-harmonic generation (SHG) has been observed in optical fibres.<sup>1</sup> The most significant part of this observation is that the amount of SH light is very small to begin with (due to the amorphous nature of glass) but with time, as the fibre is illuminated with intense laser light at  $\lambda = 1.06 \mu\text{m}$ , the SH signal grows. What is believed to happen during this 'preparation stage' is that a periodic  $\chi^{(3)}$  nonlinearity is established in the fibre. The periodicity is important for phase-matching. A good review of SHG in optical fibres can be found in Reference 2. Microscopically what is going on during the preparation is still very unclear. In this letter it is reported that for optical fibres which show efficient SHG, occasionally the third-harmonic (TH) signal can grow exponentially with time at the expense of the SH signal.

Our experiments were performed with a Q-switched (1200 Hz) and mode-locked (76 MHz) Nd: YAG laser (Quantronix 416). Peak powers between 1 and 20kW were launched into the optical fibre. The fibre used had a core diameter of  $d \approx 8.3 \mu\text{m}$ ,  $\Delta \approx 0.0037$  and a 1.3 m% phosphorous content in the core and cladding. The main dopant in the core was  $\text{GeO}_2$ .

Optical fibre lengths of 5-30 cm were used. The output of the fibre was directed into a monochromator (Jarrel-Ash, model 82-499) in conjunction with an optical multichannel analyser (OMA) (EG&G, model 1460 and a detector 1430 with an intensified linear photodiode array). A typical spectrum is displayed in Fig. 1. To exclude any contribution from the strong infra-red (IR) light at  $\lambda = 1.064 \mu\text{m}$  to the visible and UV signals, two BG18 Schott glass filters were mounted in front of the input slit of the monochromator. Apart from blocking the IR light, the filters each had  $\approx 70\%$  transmission for  $\lambda = 0.532 \mu\text{m}$  (2ω) and  $\approx 20\%$  transmission for  $\lambda = 0.355 \mu\text{m}$  (3ω). A spectrum such as the one in Fig. 1 was recorded with an integration time of 2 s, a sensitivity of  $\approx 15$  photons/count and a spectral resolution of  $\approx 6 \text{ \AA}$ . The peak around 420 nm is believed to be due to fluorescence from the TH signal; this peak has been observed by others.<sup>3</sup>




Fig. 1 Spectrum from 30 cm-long unprepared piece of fibre, illuminated with light at  $\lambda = 1.064 \mu\text{m}$ , 10kW peak power

U. ÖSTERBERG

Frank J. Seiler Research Laboratory  
US Air Force Academy, CO 80840-6328, USA

10th November 1989

For most fibres the SH signal started to grow immediately, and very soon was much larger than the TH signal. The growth rates for the SH signals for different lengths of this optical fibre can be seen in Fig. 1 of Reference 4. For approximately 10% of the fibres investigated, the TH signal started to grow exponentially with time (Fig. 2).




Fig. 2 Third- and second-harmonic signals as function of illumination time for 3 cm-long piece of fibre

The growth rate for the TH signal was the same as for the SH signal, in fibres where the SH signal grew and not the TH signal. However, in the cases when the TH signal grew exponentially with time the SH signal grew linearly, and when the TH signal saturated the SH signal started to grow exponentially with time. The saturation level for the TH signal was comparable to the saturation level for the SH signal for this particular length of fibre.

The TH signal grew to approximately 60 times its original value. This increase can be due to growth of the real part of the third-order susceptibility, an increase of the coherence length or both. If the increase could be solely accounted for by an increase in the third-order susceptibility, an increase of a factor of 11 in the spectral width of the fundamental beam should be detected, assuming  $\chi^{(3)}(\omega, \omega, -\omega, \omega) \approx \chi^{(3)}(\omega, \omega, \omega, \omega)$ . Improving our spectral resolution to  $\leq 1 \text{ \AA}$  we could detect no change in spectral width of the fundamental beam before and after the growth of the TH signal. Obviously, more measurements have to be performed before it is known what causes the TH signal to grow.

In conclusion, we report a 60 times increase of the TH signal with time in an optical glass fibre.

### References

- ÖSTERBERG, U., and MANGULIS, W.: 'Dye laser pumped by Nd: YAG laser pulses frequency-doubled in a glass optical fiber', *Opt. Lett.*, 1986, 11, p. 53
- STOLEN, R. H.: 'Second harmonic generation in optical fibers', Proc. Erice summer school in nonlinear optics, 17th-28th July 1989
- LAWANDY, N. (Brown University): private communication; AMOSOV, A. V., and PETROVSKII, G. T.: *Sov. Phys. Dokl.*, 1983, 28, (24-25)
- BATDORF, B., KRAUTSCHIK, C., ÖSTERBERG, U., STEGEVAN, G. I., LEITCH, J. W., ROTGE, J. R., and WORSE, T. F.: 'Study of the length dependence of frequency-doubled light in optical fibers', *Opt. Commun.*, accepted for publication

90 03 26 098